

Biological Agent Reference Sheet (BARS)

This content of this document is for Emory University USE ONLY.

The information and contents of this Biological Agent Reference Sheet (including all text and graphics), whether available in print or electronic format (including any digital format, e-mail transmissions, or download from the website), shall be known hereinafter as "Reference Sheet Content". The Reference Sheet Content is provided as a courtesy and is not intended as a sole source of guidance in the evaluation of Biological Agents. The Reference Sheet Content is not intended to substitute for medical advice, medical care, diagnosis or treatment obtained from a physician or health care provider. Please seek the advice of a physician or other qualified health provider with any questions you may have regarding a medical condition. Do not rely on the Reference Sheet Content for diagnosis, treatment, or medical advice. This Reference Sheet Content is for informational purposes and does not provide individualized medical care or treatment. No endorsement of any specific tests, products, or procedures is made by Reference Sheet Content or affiliated party, member, agent or employee of the Emory University Environmental Health and Safety Office.

BIOLOGICAL AGENT REFERENCE SHEET

Babesia spp.

CHARACTERISTICS	
<i>Morphology</i>	Microscopic (1-2.5 µm) protozoan parasites that infect red blood cells and are spread by certain ticks.
<i>Growth Conditions</i>	<i>Babesia</i> species are cultured in hamster erythrocytes and cell culture with fetal calf serum, RPMI 1640 medium, and antibiotics.

HEALTH HAZARDS	
<i>Host Range</i>	Rodents (mainly White-Footed Mouse), <i>Ixodes scapularis</i> ticks, and humans.
<i>Modes of Transmission</i>	Bite by infected tick; blood transfusion with contaminated blood; infected mother to baby; needle stick; open wound.
<i>Signs and Symptoms</i>	Can be asymptomatic. Common symptoms include: <ul style="list-style-type: none"> • Flu-like symptoms – fever, chills, sweats • Malaise, arthralgia, headache • Gastrointestinal symptoms – anorexia and nausea, abdominal pain, vomiting • Dark urine • Less common: cough, sore throat, depression, photophobia, red eyes • Mild enlargement of spleen, mild enlargement of liver, or jaundice may occur in some patients • Persons without a spleen can develop severe disease, even death
<i>Infectious Dose</i>	Unknown.
<i>Incubation Period</i>	1-6 weeks following tick bite; 1-9+ weeks following transfusion

MEDICAL PRECAUTIONS / TREATMENT	
<i>Prophylaxis</i>	Apply repellents with DEET to skin and clothing; avoid areas with overgrown grasses or brush; lessen amount of exposed skin; shower soon after being outdoors; check body for ticks; intervene before blood donation if donor has ever been infected.
<i>Vaccines</i>	None available.
<i>Treatment</i>	Babesiosis is usually treated for at least 7-10 days with a combination of two medications atovaquone PLUS azithromycin; OR clindamycin PLUS quinine.
<i>Surveillance</i>	Unknown.
<i>Emory Requirements</i>	Report any exposure.

LABORATORY HAZARDS	
<i>Laboratory Acquired Infections (LAIs)</i>	None reported.
<i>Sources</i>	<i>BMBL: 5th Edition</i>

SUPPLEMENTAL REFERENCES	
<i>CDC</i>	http://www.cdc.gov/parasites/babesiosis/
<i>BMBL: 5th Edition</i>	http://www.cdc.gov/OD/ohs/biosfty/bmb15/BMBL_5th_Edition.pdf

CONTAINMENT REQUIREMENTS	
<i>BSL-2</i>	For activities with infective stages of the parasite. PPE in conjunction with a BSC are required when working with cultures, tissue homogenates, or blood containing parasites.
<i>ABSL-2+</i>	For activities with infective stages of the parasite.

SPILL PROCEDURES	
<i>Small</i>	Notify others working in the lab. Allow aerosols to settle. Don appropriate PPE. Cover area of the spill with paper towels and apply an EPA registered disinfectant, working from the perimeter towards the center. Allow 30 minutes of contact time before disposal and cleanup of spill materials.
<i>Large</i>	For assistance, contact Emory's Biosafety Officer (404-727-8863), the EHSO Office (404-727-5922), or 404-727-6111 from a campus phone, or the Spill Response Team (404-727-2888).

EXPOSURE PROCEDURES		
<i>Mucous membrane</i>	Flush eyes, mouth or nose for 15 minutes at eyewash station.	
<i>Other Exposure</i>	Wash area with soap and water for 15 minutes.	
<i>Reporting</i>	Immediately report incident to supervisor, complete an employee incident report in PeopleSoft.	
<i>Medical Follow up</i>	<i>7am-4pm (OIM):</i> EUH (404-686-7941) EUHM (404-686-7106) WW (404-728-6431)	<i>After Hours:</i> OIM NP On Call 404-686-5500 PIC# 50464
	<i>Needle Stick (OIM):</i> EUH (404-686-8587) EUHM (404-686-2352)	<i>Yerkes:</i> Maureen Thompson Office (404-727-8012) Cell (404-275-0963)

VIABILITY	
<i>Disinfection</i>	Unknown.
<i>Inactivation</i>	Unknown.
<i>Survival Outside Host</i>	Unknown.

PERSONAL PROTECTIVE EQUIPMENT (PPE)	
<i>Minimum PPE Requirements</i>	At minimum, personnel are required to don gloves, closed toed shoes, lab coat, and appropriate face and eye protection prior to working with <i>Babesia</i> species. Additional PPE may be required depending on lab specific SOPs.
<i>Additional Precautions</i>	All procedures that may produce aerosols, or involve high concentrations or large volumes should be conducted in a biological safety cabinet (BSC). The use of needles, syringes, and other sharp objects should be strictly limited. Additional precautions should be considered with work involving animals or large scale activities.